



International Baccalaureate®  
Baccalauréat International  
Bachillerato Internacional

Diploma Programme

---

# Mathematics: applications and interpretation formula booklet

For use during the course and in the examinations  
First examinations 2021

Version 1.0

# Contents

## Prior learning

---

SL and HL	2
HL only	2

## Topic 1: Number and algebra

---

SL and HL	3
HL only	4

## Topic 2: Functions

---

SL and HL	5
HL only	5

## Topic 3: Geometry and trigonometry

---

SL and HL	6
HL only	7

## Topic 4: Statistics and probability

---

SL and HL	9
HL only	10

## Topic 5: Calculus

---

SL and HL	11
HL only	11

## Prior learning – SL and HL

Area of a parallelogram	$A = bh$ , where $b$ is the base, $h$ is the height
Area of a triangle	$A = \frac{1}{2}(bh)$ , where $b$ is the base, $h$ is the height
Area of a trapezoid	$A = \frac{1}{2}(a + b)h$ , where $a$ and $b$ are the parallel sides, $h$ is the height
Area of a circle	$A = \pi r^2$ , where $r$ is the radius
Circumference of a circle	$C = 2\pi r$ , where $r$ is the radius
Volume of a cuboid	$V = lwh$ , where $l$ is the length, $w$ is the width, $h$ is the height
Volume of a cylinder	$V = \pi r^2 h$ , where $r$ is the radius, $h$ is the height
Volume of prism	$V = Ah$ , where $A$ is the area of cross-section, $h$ is the height
Area of the curved surface of a cylinder	$A = 2\pi r h$ , where $r$ is the radius, $h$ is the height
Distance between two points $(x_1, y_1)$ and $(x_2, y_2)$	$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$
Coordinates of the midpoint of a line segment with endpoints $(x_1, y_1)$ and $(x_2, y_2)$	$\left( \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

## Prior learning – HL only

Solutions of a quadratic equation	The solutions of $ax^2 + bx + c = 0$ are $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ , $a \neq 0$
-----------------------------------	--

## Topic 1: Number and algebra – SL and HL

<b>SL 1.2</b>	<p>The <math>n</math>th term of an arithmetic sequence</p> <p>The sum of <math>n</math> terms of an arithmetic sequence</p>	$u_n = u_1 + (n - 1)d$ $S_n = \frac{n}{2}(2u_1 + (n - 1)d); S_n = \frac{n}{2}(u_1 + u_n)$
<b>SL 1.3</b>	<p>The <math>n</math>th term of a geometric sequence</p> <p>The sum of <math>n</math> terms of a finite geometric sequence</p>	$u_n = u_1 r^{n-1}$ $S_n = \frac{u_1(r^n - 1)}{r - 1} = \frac{u_1(1 - r^n)}{1 - r}, r \neq 1$
<b>SL 1.4</b>	Compound interest	$FV = PV \times \left(1 + \frac{r}{100k}\right)^{kn}$ <p>where <math>FV</math> is the future value,  <math>PV</math> is the present value, <math>n</math> is the number of years,  <math>k</math> is the number of compounding periods per year,  <math>r\%</math> is the nominal annual rate of interest</p>
<b>SL 1.5</b>	Exponents and logarithms	$a^x = b \Leftrightarrow x = \log_a b, \text{ where } a > 0, b > 0, a \neq 1$
<b>SL 1.6</b>	Percentage error	$\varepsilon = \left  \frac{v_A - v_E}{v_E} \right  \times 100\%$ <p>where <math>v_E</math> is the exact value and <math>v_A</math> is the approximate value of <math>v</math></p>

## Topic 1: Number and algebra – HL only

<b>AHL 1.9</b>	Laws of logarithms	$\log_a xy = \log_a x + \log_a y$ $\log_a \frac{x}{y} = \log_a x - \log_a y$ $\log_a x^m = m \log_a x$ for $a, x, y > 0$
<b>AHL 1.11</b>	The sum of an infinite geometric sequence	$S_\infty = \frac{u_1}{1-r},  r  < 1$
<b>AHL 1.12</b>	Complex numbers  Discriminant	$z = a + bi$  $\Delta = b^2 - 4ac$
<b>AHL 1.13</b>	Modulus-argument (polar) and exponential (Euler) form	$z = r(\cos \theta + i \sin \theta) = re^{i\theta} = r \operatorname{cis} \theta$
<b>AHL 1.14</b>	Determinant of a $2 \times 2$ matrix  Inverse of a $2 \times 2$ matrix	$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow \det A =  A  = ad - bc$  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow A^{-1} = \frac{1}{\det A} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}, ad \neq bc$
<b>AHL 1.15</b>	Power formula for a matrix	$M^n = PD^nP^{-1}$ , where $P$ is the matrix of eigenvectors and $D$ is the diagonal matrix of eigenvalues

## Topic 2: Functions – SL and HL

<b>SL 2.1</b>	Equations of a straight line  Gradient formula	$y = mx + c$ ; $ax + by + d = 0$ ; $y - y_1 = m(x - x_1)$  $m = \frac{y_2 - y_1}{x_2 - x_1}$
<b>SL 2.5</b>	Axis of symmetry of the graph of a quadratic function	$f(x) = ax^2 + bx + c \Rightarrow$ axis of symmetry is $x = -\frac{b}{2a}$

## Topic 2: Functions – HL only

<b>AHL 2.9</b>	Logistic function	$f(x) = \frac{L}{1 + Ce^{-kx}}$ , $L, k, C > 0$
--------------------	-------------------	---

## Topic 3: Geometry and trigonometry – SL and HL

<b>SL 3.1</b>	<p>Distance between two points <math>(x_1, y_1, z_1)</math> and <math>(x_2, y_2, z_2)</math></p> <p>Coordinates of the midpoint of a line segment with endpoints <math>(x_1, y_1, z_1)</math> and <math>(x_2, y_2, z_2)</math></p> <p>Volume of a right-pyramid</p> <p>Volume of a right cone</p> <p>Area of the curved surface of a cone</p> <p>Volume of a sphere</p> <p>Surface area of a sphere</p>	$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2}$ $\left( \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right)$ $V = \frac{1}{3} Ah, \text{ where } A \text{ is the area of the base, } h \text{ is the height}$ $V = \frac{1}{3} \pi r^2 h, \text{ where } r \text{ is the radius, } h \text{ is the height}$ $A = \pi r l, \text{ where } r \text{ is the radius, } l \text{ is the slant height}$ $V = \frac{4}{3} \pi r^3, \text{ where } r \text{ is the radius}$ $A = 4\pi r^2, \text{ where } r \text{ is the radius}$
<b>SL 3.2</b>	<p>Sine rule</p> <p>Cosine rule</p> <p>Area of a triangle</p>	$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ $c^2 = a^2 + b^2 - 2ab \cos C; \cos C = \frac{a^2 + b^2 - c^2}{2ab}$ $A = \frac{1}{2} ab \sin C$
<b>SL 3.4</b>	<p>Length of an arc</p> <p>Area of a sector</p>	$l = \frac{\theta}{360} \times 2\pi r, \text{ where } \theta \text{ is the angle measured in degrees, } r \text{ is the radius}$ $A = \frac{\theta}{360} \times \pi r^2, \text{ where } \theta \text{ is the angle measured in degrees, } r \text{ is the radius}$

## Topic 3: Geometry and trigonometry – HL only

<b>AHL 3.7</b>	Length of an arc  Area of a sector	$l = r\theta$ , where $r$ is the radius, $\theta$ is the angle measured in radians  $A = \frac{1}{2}r^2\theta$ , where $r$ is the radius, $\theta$ is the angle measured in radians
<b>AHL 3.8</b>	Identities	$\cos^2 \theta + \sin^2 \theta = 1$  $\tan \theta = \frac{\sin \theta}{\cos \theta}$
<b>AHL 3.9</b>	Transformation matrices	$\begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix}$ , reflection in the line $y = (\tan \theta)x$  $\begin{pmatrix} k & 0 \\ 0 & 1 \end{pmatrix}$ , horizontal stretch / stretch parallel to $x$ -axis with a scale factor of $k$  $\begin{pmatrix} 1 & 0 \\ 0 & k \end{pmatrix}$ , vertical stretch / stretch parallel to $y$ -axis with a scale factor of $k$  $\begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}$ , enlargement, with a scale factor of $k$ , centre $(0, 0)$  $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ , anticlockwise/counter-clockwise rotation of angle $\theta$ about the origin ( $\theta > 0$ )  $\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$ , clockwise rotation of angle $\theta$ about the origin ( $\theta > 0$ )



<b>AHL 3.10</b>	Magnitude of a vector	$ \mathbf{v}  = \sqrt{v_1^2 + v_2^2 + v_3^2}$ , where $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$
<b>AHL 3.11</b>	Vector equation of a line  Parametric form of the equation of a line	$\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$  $x = x_0 + \lambda l, y = y_0 + \lambda m, z = z_0 + \lambda n$
<b>AHL 3.13</b>	Scalar product  Angle between two vectors  Vector product  Area of a parallelogram	$\mathbf{v} \cdot \mathbf{w} = v_1 w_1 + v_2 w_2 + v_3 w_3$ , where $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$ , $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$ $\mathbf{v} \cdot \mathbf{w} =  \mathbf{v}   \mathbf{w}  \cos \theta$ , where $\theta$ is the angle between $\mathbf{v}$ and $\mathbf{w}$ $\cos \theta = \frac{v_1 w_1 + v_2 w_2 + v_3 w_3}{ \mathbf{v}   \mathbf{w} }$ $\mathbf{v} \times \mathbf{w} = \begin{pmatrix} v_2 w_3 - v_3 w_2 \\ v_3 w_1 - v_1 w_3 \\ v_1 w_2 - v_2 w_1 \end{pmatrix}$ , where $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$ , $\mathbf{w} = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$ $ \mathbf{v} \times \mathbf{w}  =  \mathbf{v}   \mathbf{w}  \sin \theta$ , where $\theta$ is the angle between $\mathbf{v}$ and $\mathbf{w}$ $A =  \mathbf{v} \times \mathbf{w} $ where $\mathbf{v}$ and $\mathbf{w}$ form two adjacent sides of a parallelogram

## Topic 4: Statistics and probability – SL and HL

<b>SL 4.2</b>	Interquartile range	$IQR = Q_3 - Q_1$
<b>SL 4.3</b>	Mean, $\bar{x}$ , of a set of data	$\bar{x} = \frac{\sum_{i=1}^k f_i x_i}{n}$ , where $n = \sum_{i=1}^k f_i$
<b>SL 4.5</b>	Probability of an event $A$	$P(A) = \frac{n(A)}{n(U)}$
	Complementary events	$P(A) + P(A') = 1$
<b>SL 4.6</b>	Combined events	$P(A \cup B) = P(A) + P(B) - P(A \cap B)$
	Mutually exclusive events	$P(A \cup B) = P(A) + P(B)$
	Conditional probability	$P(A B) = \frac{P(A \cap B)}{P(B)}$
	Independent events	$P(A \cap B) = P(A) P(B)$
<b>SL 4.7</b>	Expected value of a discrete random variable $X$	$E(X) = \sum x P(X = x)$
<b>SL 4.8</b>	Binomial distribution $X \sim B(n, p)$	
	Mean	$E(X) = np$
	Variance	$\text{Var}(X) = np(1 - p)$

## Topic 4: Statistics and probability – HL only

<b>AHL 4.14</b>	<p>Linear transformation of a single random variable</p> <p>Linear combinations of <math>n</math> independent random variables, <math>X_1, X_2, \dots, X_n</math></p> <p>Sample statistics</p> <p>Unbiased estimate of population variance <math>s_{n-1}^2</math></p>	$E(aX + b) = aE(X) + b$ $\text{Var}(aX + b) = a^2 \text{Var}(X)$ $E(a_1X_1 \pm a_2X_2 \pm \dots \pm a_nX_n) = a_1E(X_1) \pm a_2E(X_2) \pm \dots \pm a_nE(X_n)$ $\text{Var}(a_1X_1 \pm a_2X_2 \pm \dots \pm a_nX_n)$ $= a_1^2 \text{Var}(X_1) + a_2^2 \text{Var}(X_2) + \dots + a_n^2 \text{Var}(X_n)$ $s_{n-1}^2 = \frac{n}{n-1} s_n^2$
<b>AHL 4.17</b>	<p>Poisson distribution <math>X \sim \text{Po}(m)</math></p> <p>Mean</p> <p>Variance</p>	$E(X) = m$ $\text{Var}(X) = m$
<b>AHL 4.19</b>	Transition matrices	$T^n s_0 = s_n$ , where $s_0$ is the initial state

## Topic 5: Calculus – SL and HL

<b>SL 5.3</b>	Derivative of $x^n$	$f(x) = x^n \Rightarrow f'(x) = nx^{n-1}$
<b>SL 5.5</b>	Integral of $x^n$  Area of region enclosed by a curve $y = f(x)$ and the $x$ -axis, where $f(x) > 0$	$\int x^n dx = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1$  $A = \int_a^b y dx$
<b>SL 5.8</b>	The trapezoidal rule	$\int_a^b y dx \approx \frac{1}{2}h((y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1}))$ , where $h = \frac{b-a}{n}$

## Topic 5: Calculus – HL only

<b>AHL 5.9</b>	Derivative of $\sin x$	$f(x) = \sin x \Rightarrow f'(x) = \cos x$
	Derivative of $\cos x$	$f(x) = \cos x \Rightarrow f'(x) = -\sin x$
	Derivative of $\tan x$	$f(x) = \tan x \Rightarrow f'(x) = \frac{1}{\cos^2 x}$
	Derivative of $e^x$	$f(x) = e^x \Rightarrow f'(x) = e^x$
	Derivative of $\ln x$	$f(x) = \ln x \Rightarrow f'(x) = \frac{1}{x}$
	Chain rule	$y = g(u), \text{ where } u = f(x) \Rightarrow \frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$
	Product rule	$y = uv \Rightarrow \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$
Quotient rule	$y = \frac{u}{v} \Rightarrow \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$	

<b>AHL 5.11</b>	Standard integrals	$\int \frac{1}{x} dx = \ln x  + C$ $\int \sin x dx = -\cos x + C$ $\int \cos x dx = \sin x + C$ $\int \frac{1}{\cos^2 x} = \tan x + C$ $\int e^x dx = e^x + C$
<b>AHL 5.12</b>	Area of region enclosed by a curve and $x$ or $y$ -axes  Volume of revolution about $x$ or $y$ -axes	$A = \int_a^b  y  dx \text{ or } A = \int_a^b  x  dy$ $V = \int_a^b \pi y^2 dx \text{ or } V = \int_a^b \pi x^2 dy$
<b>AHL 5.13</b>	Acceleration  Distance travelled from $t_1$ to $t_2$  Displacement from $t_1$ to $t_2$	$a = \frac{dv}{dt} = \frac{d^2s}{dt^2} = v \frac{dv}{ds}$ $\text{distance} = \int_{t_1}^{t_2}  v(t)  dt$ $\text{displacement} = \int_{t_1}^{t_2} v(t) dt$
<b>AHL 5.16</b>	Euler's method  Euler's method for coupled systems	$y_{n+1} = y_n + h \times f(x_n, y_n); x_{n+1} = x_n + h, \text{ where } h \text{ is a constant (step length)}$ $x_{n+1} = x_n + h \times f_1(x_n, y_n, t_n)$ $y_{n+1} = y_n + h \times f_2(x_n, y_n, t_n)$ $t_{n+1} = t_n + h$ <p>where <math>h</math> is a constant (step length)</p>
<b>AHL 5.17</b>	Exact solution for coupled linear differential equations	$x = Ae^{\lambda_1 t} p_1 + Be^{\lambda_2 t} p_2$